



BioTrace Ltd
 Unit 4 - 110 Mays Road
 1061 Onehunga, Auckland
 Tél.:09 570 5923

Patient: John Smith

Date: 2018/04/09
 Male 62 years
 Date of Birth: yyyy/mm/dd
 Blood group: A
 Weight: 76 Kg
 Size: 1m 76

Mineral Test Report

		Result	Normal	Low-	Low	Normal	OK	Normal+	High	High+
Calcium	Ca	511.0	279.0	598.0						
Magnesium	Mg	27.2	30.5	75.7						
Phosphorus	P	131.3	144.0	199.0						
Silicon	Si	11.9	15.0	31.0						
Sodium	Na	49.4	21.0	89.0						
Potassium	K	16.0	9.0	39.0						
Copper	Cu	17.4	11.0	28.0						
Zinc	Zn	135.8	125.0	155.0						
Iron	Fe	9.4	5.0	15.0						
Manganese	Mn	0.42	0.31	0.75						
Chromium	Cr	0.70	0.82	1.25						
Vanadium	V	0.024	0.009	0.083						
Boron	B	2.28	0.84	2.87						
Cobalt	Co	0.031	0.025	0.045						
Molybdenum	Mo	0.042	0.035	0.085						
Iodine	I	0.42	0.32	0.59						
Lithium	Li	0.067	0.052	0.120						
Germanium	Ge	0.019	0.003	0.028						
Selenium	Se	0.65	0.95	1.77						
Sulphur	S	48.3	48.1	52.0						

Mineral Balance





Heavy Metal Test Report

	Result	Normal	High -	High +	Excess
Aluminium (Al)	0.01225				
Antimony (Sb)	0.00235				
Silver (Ag)	0.01363				
Arsenic (As)	0.00469				
Barium (Ba)	0.00611				
Beryllium (Be)	0.00562				
Bismuth (Bi)	0.00763				
Cadmium (Cd)	0.01046				
Mercury (Hg)	0.01613				
Nickel (Ni)	0.00377				
Platinum (Pt)	0.00236				
Lead (Pb)	0.00598				
Thallium (Tl)	0.00148				
Thorium (Th)	0.00091				

Heavy Metals Intoxication

Overall Intoxication

unsatisfactory: 82%



Blockage suspicion for heavy metals elimination; possibly from lack of sulfur conjugation.

unsatisfactory: 75%



Ratios

	Ratios	Normal	Low	OK	Haut	Deficiency	Excess
Ca/Mg	18.81	7.84 - 18.25				Mg	
Ca/P	3.89	1.64 - 4.15				P	
K/Na	0.32	0.45 - 0.75					
Cu/Zn	0.13	0.11 - 0.17					

Oxidative Stress


Oxidative Aggression

good: 60%



Oxidative Protection

good: 70%





BioTrace Ltd
Unit 4 - 110 Mays Road
1061 Onehunga, Auckland
Tél.:09 570 5923

Patient: John Smith

Date: 2018/04/09
Male 62 years
Date of Birth: yyyy/mm/dd
Blood group: A
Weight: 76 Kg
Size: 1m 76

Comments

Methodology: Spectrophotometry of palmar dermis
Comments:

Caution! These values are not for diagnostic purposes ; these are only an interprétation of corrélations between minerals & oligo-elements tested with Oligoscan. These relationships have been widely documented throughout the scientific literature on micronutrients and ortho-molecular medicine. The Oligoscan test is only a functional element of balance in the body.



OligoScan Report

John Smith

9th April 2018

Mineral Deficiencies

Below are the minerals appearing deficient or low on John's OligoScan.

Magnesium (Mg)

Functions

- Mg is required for the activation and/or structure of more enzymes than any other mineral. Many Mg dependent enzymes cannot use a different mineral to replace Mg
- Regulation of cell membranes: permeability, muscle contraction, nerve impulse conduction and antagonism to Calcium (Ca)
- Required for bone formation
- Detoxification: required for optimal liver activity
- Metabolic: required for energy production, glucose and fat metabolism, and for protein synthesis
- Immune: Mg operates at several levels in immune mechanisms such as inflammatory reactions, allergies and boosting white blood cells

Deficiency: Mg is blocked by Cadmium, Copper, Aluminium, Mercury and Fluoride and is used up rapidly during times of stress as adrenaline keeps Mg suspended in the blood, increasing Mg loss.

Dietary Sources

Green leafy vegetables esp. spinach, kale, and silver beet, vegetables esp. broccoli, corn, and avocado, fruits (dried – apple, figs, apricots, dates, currants, raisins, sultanas, prunes; fresh – passion fruit, blackberry, raspberry, banana), nuts, seeds, tahini, whole-grains, dried seaweed, cocoa powder, and dark chocolate.

Recommended Mg Supplement: BioTrace Coral Cal-Mag

Phosphorus (P)

Functions

- Bone structure: 80-85% of P is located in the teeth and bones
- Energy/nervous/muscular: ATP and ADP for energy and muscle contraction
- Component of myelin sheath which covers nerve cells
- Cell membrane: as phospholipids
- Genetic reactions: in DNA and RNA
- Buffering agent to maintain osmotic pressure
- Digestive: regulates absorption of Calcium, and a variety of trace elements
- Immune: ATP for leukocytes
- Detoxification: in liver via ATP

Deficiency: P may be blocked by Lead or Aluminium. Low P may also be an indicator for insufficient protein synthesis due to either low dietary protein or low digestive enzymes.

Dietary Sources

P is found in red meat, fish and shellfish, natural unsweetened yoghurt, eggs, fruit (dried – apricots, currants, dates, figs, sultanas, prunes; fresh – banana, all berries, peaches, plums, nectarines, grapes), legumes & beans, tofu, seeds (sunflower, sesame), tahini, and vegetables (mushrooms, sweet corn, sundried tomato, peas).

Recommended P Supplementation: PRL Greens helps balance P and Ca levels, as well as body pH.

Silicon (Si)

Functions

- The presence of Si is necessary for the biosynthesis of collagen, elastin and hyaluronic acid. Therefore, Si plays an important role in connective tissue health, specifically cartilage, bone, skin, hair, and nails
- Si is needed for the synthesis of all membranes in the body including the fascia, meninges, periosteum, and intestinal support membranes
- Si is important for the cerebellum, a part of the brain associated with maintaining balance in the body and mind
- A low Si state creates a situation where there is insufficient storage for minerals

Deficiency: Si is depleted by *Aluminium* and is required to help antagonise *Aluminium* out of cells.

Dietary Sources

Horsetail herb, whole-grains (oats, brown rice, barley, millet, corn, rye), flaxseeds, black tea, banana, raisins, vegetables (green beans, carrot, cucumber (skin), chickweed, alfalfa, artichoke, red beets, asparagus, potatoes, stems of green vegetables), and kelp.

Recommended Si Supplement: PRL Greens

Chromium (Cr)

Functions

- Cr is required for the body's production of GTF (glucose tolerance factor). This has a role in regulating the pancreatic hormones responsible for controlling blood glucose
- This GTF complex also affects lipid metabolism and reduces cholesterol
- Low cellular levels of bio-available Cr are associated with carbohydrate intolerance and fluctuations in insulin production and blood sugar levels that may contribute to weight gain, energy imbalances, sugar cravings, and mood disturbances

Deficiency: Magnesium, Zinc and Manganese are synergistic minerals for glucose metabolism. Low Cr may indicate low stomach acid (HCL). Sugars, starch, fruit, fruit juices and stress and the resultant cortisol and insulin secretion can deplete chromium.

Dietary Sources

Cr is in brewer's yeast, organic wholegrain cereals, seafood, oysters, liver, chicken, beef, potatoes, apples, bananas, spinach, butter and thyme. Refined foods and intensive farming reduces the presence of Cr in food. The consumption of sugar and refined flour tends to accelerate its decline.

Recommended Cr Supplement: BIO Elemental Chromium

Selenium (Se)

Functions

- Various studies on this major antioxidant have shown its protective and preventive role in many diseases
- Se is a coenzyme for glutathione synthesis, which protects the cell membrane and the nucleus against oxidation when under attack by free radicals
- It protects against the toxic effects of heavy metals (*Arsenic*, *Hg*, *Cd* and *Pb*), alcohol, tobacco smoke and various atmospheric pollutions

- Combined with Vitamin E, Se induces the formation of antibodies and prevents oxidative damage to chromosomes. Thus, together they *boost immunity* by enhancing the activation and proliferation of B lymphocytes and by strengthening the functioning of T cells
- Se protects the cardiovascular system by controlling the optimum amount of red blood cells, standardising platelet aggregation and accelerating fat metabolism. It is also a regulatory agent in blood pressure and heart rate

Deficiency: Mercury is a major antagonist for Se, as are Cadmium, Arsenic and Lead. Adequate Se levels are needed to help detoxify heavy metals.

Dietary Sources

Brazil nuts, eggs, mushrooms, red meat, chicken, whole grains (oat bran, oats, wheat bran & germ, rye flour, brown rice), mustard powder, sesame seeds & tahini, and all fish.

Recommended Se Supplement: [Bio El Selenium](#)

Sulphur (S)

Functions:

- S is involved in the composition of different amino acids.
- It is involved in many metabolic functions particularly in connective tissue and is often recommended in arthritis or osteoarthritis
- It has anti-allergic properties (like manganese) and can be very useful in skin diseases, eczema and dermatitis
- Required for detoxification and excretion pathways. It binds toxins to amino acids for excretion

Deficiency: S is an important cofactor for chelation of toxic metals from the body. Low S on OligoScan can indicate impairments in detoxification pathways. Abnormally high levels of Calcium, Magnesium, Phosphorous or Zinc may indicate a Sulphur blockade.

Dietary Sources

Eggs, meat, poultry, fish, milk, horseradish, onions, garlic, red pepper, yeast, soy, legumes, and organic whole grains

Recommended S support Supplement: [PRL NAC](#)

Mineral Excesses

Below are the minerals appearing in excess or raised on John's OligoScan. Mineral Excesses on OligoScan indicate either over-supplementation or a blockade – meaning the mineral is bio-unavailable. This can happen due to other minerals or toxic metals blocking the absorption and utilisation of the mineral in question.

Calcium (Ca)

Functions

- Ca is an essential mineral required for the function of numerous intracellular and extracellular processes including muscle contraction, nerve conduction, beating of the heart, hormone release, blood coagulation, energy production and maintenance of immune function
- Ca is regulated by the parathyroid hormone (PTH), calcitonin and vitamin D
- It is involved in bone and teeth mineralisation
- It plays a role in intracellular signalling and is involved in the regulation of many enzymes responsible for fat digestion and protein metabolism
- Ca absorption is impaired in achlorhydria (low stomach acid), intestinal inflammation and many malabsorptive disorders
- Ca deficiency may lead to muscle pain and spasms, bone pain, osteoporosis and increased risk of fractures, tooth decay, altered heart rate, and ambulatory developmental delay in children

Excess on OligoScan: Ca is blocked by Cadmium and Lead, and balanced by Magnesium, Phosphorous, Boron, Silicon and Molybdenum. When two or more of the bone minerals are out of balance tissue acidity (low pH) is

suspected. The blood sequesters Ca from bone in an effort to balance the blood's pH. This in turn affects the other bone mineral levels.

Dietary Sources

Unpasteurised milk, dairy products, fish with bones, especially salmon and sardines, tofu, broccoli, collard greens, mustard greens, bok choy, clams and black strap molasses.

Recommended alkalising supplement: [PRL Greens](#)

Boron (B)

Functions

- Adequate levels of dietary boron are thought to play a significant role in calcium and bone metabolism, to help support healthy bone mineralisation and strong, cavity resistant teeth
- Boron has been linked to Vitamin D deficiency, low testosterone and oestrogen metabolism
- Boron may support healthy brain function and hormonal balance

Excess on OligoScan: If not supplementing, excess Boron on Oligoscan can indicate a blockade. When coupled with other 'bone mineral' deficiencies such as Calcium, Magnesium, Phosphorous and/or Silicon, low pH (acidity) is suspected.

Dietary Sources

Highest concentrations of B are found in avocado, peanut butter, peanuts, prune juice, grape juice, cocoa powder, wine, pecans, and raisins. Other sources are legumes, coffee, milk, apples and potatoes.

Recommended alkalising supplement: [PRL Greens](#)

Germanium (Ge)

Functions

- The function of Ge in humans is not well known but it has been linked to immune function and may act as an immuno-stimulant according to preliminary research studies
- Ge is linked to selenium blockades, often caused by overt or hidden mercury toxicity and/or zinc deficiency

Excess on OligoScan: This is often a sign of Mercury toxicity making the Ge bio-unavailable. Mercury antagonises Ge, thus Ge deficiency may be associated with Mercury toxicity symptoms.

Dietary Sources

Broccoli, celery, garlic, shitake mushrooms, milk, chlorella, onions, pearl barley, rhubarb sauerkraut, and tomato juice.

Mobilised Toxic Metals

Toxic metals can be harmful even in small amounts and have no known function in the body. If toxic metals enter the body and accumulate faster than the body can detoxify them, a gradual build-up will occur. Therefore, high-concentration exposure is not necessary to produce a state of toxicity as toxic metals accumulate in bodily tissues over time.

The accumulation of toxic metals can have adverse effects, even in small amounts. This is due to the manner of interaction within the body. Signs and symptoms may be nonspecific at first, and symptoms will be unique to the individual. Common symptoms may include digestive upsets, fatigue and/or a range of sensitivities. Long-term signs and symptoms of dysfunction or disease may develop. This will depend on the element concerned and the significance of the toxicity, as well as the mineral deficiencies and/or excesses exhibited by the individual.

Toxic Metals elevated for John are:

Aluminium (Al)

Sources

- Tap water – city treatment plants produce drinking water that use Al sulphate to eliminate microorganisms and organic matter from water, this is left in our drinking water
- Kitchen utensils like pots or certain packaging (rolls of Al foil or trays) release Al food. Canned food and soda cans also contain Al
- Food additives – dyes (E173), anticoagulants (E520 - E521-E522-E523) used in food made from egg whites, in pastries (E541, E555), in dried food powder (milk, coffee and soup powders, refined industrial cheese (E554-E556-E559), and baking powder
- Table salt contains aluminium as an anti-caking agent. Processed foods containing table salt will also contain aluminium
- Cosmetics such as some toothpaste, anti-perspirant deodorant, concealer, eye-shadow, talc, and foundation may contain Al chloride
- Most vaccines contain Al hydroxide which is used as an adjuvant to stimulate an immune response to the vaccine
- Medications - many antacids contain Al hydroxide, such as Gaviscon

Signs of Excess Al

- Symptoms can include: headaches, digestive tract dysbiosis, dry skin, hypoparathyroidism, kidney and liver dysfunction, neuromuscular and neurological disorders
- Al is toxic for the brain and can affect memory and cognition. Several studies show that the risk of developing Alzheimer's disease is eight times higher in people who have an unusually high amount of Al
- Al can interfere with the absorption of Calcium, Magnesium and Phosphorus. This may lead to potential bone metabolism issues
- Al toxicity may also lead to muscle aches, anaemia, digestive disorders, impaired liver function, and renal impairment
- Inflammatory muscle disease (macrophagic myofasciitis) whose symptoms are muscle pain and fatigue with a slight fever. Muscle biopsies (cell samples) performed on these patients reveal the presence of an abnormal concentration of Al in the muscle where they received a vaccine containing Al
- Disruptive action on the immune system: primary cause of autoimmune diseases

Silver (Ag)

Silver (Ag) tends to mobilise with Aluminium thus often appears high in sync with Al. Silver usually stays in the periphery of the body and doesn't lodge in the organs or brain like Al, thus is not such a concern unless it is very high.

Sources:

- Dental amalgams along with mercury
- Jewellery industry
- Photograph developer chemicals
- Surgical implants may be coated with silver
- Supplementation with colloidal silver

Signs of Excess Ag:

- Though rare, symptoms of excess Ag often present as symptoms of Aluminium and/or Mercury toxicity

Barium (Ba)

Sources:

- Commonly seen in people after digestive or oral motor function investigations (Ba enema, Ba swallow)
- In soils and foods such as nuts, seaweed, fish, and some plants

- Tattoo ink contains some Ba based chemicals (providing colour)
- Occupational exposure – Ba is used in the manufacture of many items:
 - Alloys, lubricants, electrodes, for nickel-barium parts (ignition devices)
 - Dyes and finishes for the textile industry, pigments, inks and varnishes
 - Coloured and optical glass, ceramics, paints, paper, artificial marble, cloth, enamels, rubber, plastics
 - In the refining of Al
 - Pesticides
 - Water softeners, drilling fluids, corrosion inhibitors
 - Refining animal and vegetable oils
 - Sugar industry

Signs of Excess Ba:

- Ba can be an indicator for the presence of xenoestrogens
- Interference with Potassium metabolism
- Long term exposure to Ba can cause cardiomyopathy, heart palpitations and angina
- Abdominal cramps
- Disruptive effect on the immune system

Beryllium (Be)

Sources:

- Air pollution (burning fossil fuels), nuclear applications, volcanic ash
- Manufacture of plastics, ceramics, electronics, computer components, dental plates, x-ray tubes, sporting goods, energy saving bulbs, and steel alloys
- Present in many industries including jewellery, dental, optical, metallurgy, electronic waste recycling, and aerospace

Signs of Excess Be:

- Be is irritating, allergenic and carcinogenic (high levels of reported lung cancer)
- Acute exposure leads to symptoms similar to that seen in pneumonia and bronchitis
- Shortness of breath
- Fatigue
- Loss of appetite
- Weight loss
- Joint pain
- Night sweats
- Fever
- Progressive respiratory failure
- Symptoms of chronic Be disease include breathing difficulties, coughing, chest pain, and general weakness. Signs include enlargement of the liver, spleen, and right heart, as well as the formation of kidney stones
- Granulomas on the lung

- Disruptive action on the immune system

Bismuth (Bi)

Sources:

- Bi is used in cast iron welding manufacture and in fusible alloys such as in bird pellets and fishing sinkers with a low toxicity
- Some Bi compounds are manufactured and used in the pharmaceutical industry (antacids, anti-haemorrhoid drugs, anti-ulcer drugs)
- Bismuth oxide is used in some dental materials including hydraulic silicilate cement (HSC) which is used for pulp regeneration, capping, root canal sealing and filling
- Hair dyes and cosmetics (lipstick)
- Used as a non-toxic replacement for lead in lubricating greases, oil and acrylic paints, and glazes

Signs of Excess Bi:

- Mostly affects the kidney, liver and bladder. Skin and respiratory irritation can follow
- Lack of energy, confusion, headaches, and memory loss
- Decreased appetite, weight loss and diarrhoea
- Joint pain, skin rashes, tremors
- Formation of black deposits on the gingival (bismuth line)
- Insomnia

Cadmium (Cd)

Sources

- For smokers, the major source of Cd exposure is cigarette smoke
- For non-smokers, the major source of Cd intake is food. This is due to the fact that Cd is present in trace amounts in food products: Cd that is present in the soil is easily absorbed by vegetables. Cd is a by product found in *superphosphate fertilisers*, so it can be found in the food chain. NZ has very high use of these fertilisers
- Cd is used mainly in:
 - The rechargeable nickel-cadmium batteries, solar cells
 - Electronic appliances
 - In plastic, glass, and ceramic as pigments
 - Artists' paints
 - Coatings of metals and alloys to make them resistant to corrosion
 - Burning plastics
 - Coal fired industry
 - Galvanised water pipes
 - Mining, smelting and refining
 - Paper mills
 - Exhaust fumes from automobiles

Signs of Excess Cd:

- Fatigue is a key symptom, with trouble waking in the morning
- Difficulty concentrating
- Headaches
- Cd inhibits Magnesium and Zinc levels, as well as inhibiting glucose uptake
- Skin rashes
- An increased risk of bone fractures due to osteomalacia and osteoporosis

- Joint pain, gout, arthritis
- Liver/detoxification issues
- Hyperlipidaemia
- Muscle weakness
- Cd antagonises Calcium causing hardening and weakening of arteries and inflammation potentially leading to cardiovascular disease, arteriosclerosis, hypertension, or stroke
- Disruptive action on the immune system

Mercury (Hg)

Sources

- Dental amalgams*
- Preservative in vaccines (Thimerosal)
- Medications
 - Mercurochrome/Merbromin
 - Merthiolate – preservative in topical cream
 - Preparation H – haemorrhoid cream
 - Organomercurials – mercurial diuretics
- Congenital exposure
- Contaminated large fish and shellfish, contaminated drinking water
- Seeds and vegetables treated with mercurial fungicides, some fertilisers
- Used in the manufacture of paper, felt, chlorine, adhesives, fabric softeners, and waxes
- Gold industry
- Plastics
- Printing ink, some paints, tattoo ink
- Pesticides (organo-mercurial)
- Contact lens solution
- Neon lights, energy saving light bulbs
- Cosmetics (mascara, skin lightening agents)

*A person averaging eight amalgams in their mouth is intoxicated with 15 mcg Hg per day. In comparison, environmental Hg pollution and consumption of Hg-contaminated fish is only 2 mcg per day. In the mouth, a filling undergoes both mechanical abrasion and electrochemical corrosion. Hg is released mainly in the form of vapour and enters the body through breathing. When measuring the fumes coming out of a mouth with amalgam fillings, the value exceeds 480 times the accepted standard according to Ministry for the Environment NZ.

Signs of Excess Hg:

- Inhibits glucose transfer and the production of insulin and can be a factor in hypoglycaemia and diabetes
- Inhibits Mg and Zn levels
- Muscle tremors, muscle weakness, paralysis, convulsions, numbness and tingling in arms and legs
- Digestive problems, salivation, stomatitis, periodontitis
- Kidney damage
- Vision loss
- Hyperactivity and attention disorders in children, autism
- Fatigue, insomnia
- Neurological problems (memory loss, anxiety, depression, mood swings)
- Endocrine dysfunction, thyroid and immune dysfunction

Lead (Pb)

Sources:

- Contaminated dust and dirt is often how children are exposed to Pb
- Pb car batteries and other battery manufacture
- Mine and smelting industries

- Old water pipes and contaminated drinking water
- In the production of ceramics, glazes, PVC plastic, inks, ammunition, crystal, and fishing sinkers
- Cigarette smoke
- Food cans smouldered with Pb
- Pesticide residues
- Pb sheets used in the construction sector
- Some hair dyes (black)
- Previously used as a pigment in paints and an octane in gasoline (led to contaminated soil, water and air)
- Congenital exposure

Signs of Excess Pb:

- Saturnism (lead poisoning) refers to people with all of the indications of Pb intoxication
- Pb replaces Calcium through enzyme binding sites, especially in bone. Can lead to weak bones and osteoporosis
- Hinders activity of Magnesium, Vitamin B1 and Zinc
- Arthritis, gout, back pain
- Epilepsy
- Fatigue, insomnia
- Neurological disorders
 - Behavioural problems: hyperactivity, attention deficit disorder
 - Mental weakness, dyslexia
- Effects on bone marrow and blood – Pb blocks several enzymes needed for haemoglobin synthesis. May lead to anaemia
- Gastrointestinal problems
 - Constipation or diarrhoea
 - Metallic taste in the mouth
 - Abdominal pain or cramping
- Disrupts the immune system

Significant Mineral Ratios

The following significant mineral ratios are imbalanced:

Calcium/Magnesium Ratio (Pancreas/Insulin/Lifestyle ratio)

Increased Ca/Mg ratio

Indicates Mg deficiency. Increased insulin levels may be released leading to hypoglycaemic symptoms such as hunger, shakiness, nervousness, sweating, dizziness or light-headedness, sleepiness, confusion, anxiety, and weakness. The following may also be indicated:

- Lifestyle stress
- Eating high carbohydrate diet (leading to blood sugar stress)
- Osteoporosis
- Periodontal disease
- Hyperthyroidism or hyperparathyroidism
- Cardiovascular system problems
- Defect in Calcium absorption pathway with possible inappropriate Calcium deposition

Potassium to Sodium (K/Na) Ratio (Vitality/Kidney & Adrenals ratio)

Decreased K/Na ratio

Loss of K and/or excess retention of Na. The following may be indicated:

- Sympathetic state, characterised by acute stress and increased adrenal gland activity
- Adrenal gland hyperactivity leading to hypertension, anxiety, and cardiovascular problems
- Inflammation

- Excessive consumption of salt (processed and refined foods, or added to foods)
- Loop diuretics

Dietary Suggestions for John

- ↑ **Vegetables:** (as suggested in food sources of deficient minerals) to help increase alkalinity of the body
 - ↓ **Sugar:** sugar increases body acidity and depletes nutrients, especially Magnesium
 - ↓ **Coffee and black tea:** intake as these can affect mineral absorption, especially Magnesium, Calcium and *Iron*
 - ↑ **High quality protein:** in your diet to help enhance liver detoxification pathways
 - ↑ **Sulphur containing foods:** such as *Brassicas* (cabbage, broccoli, cauliflower, brussel sprouts, and bok choy) and *Alliums* (onions, shallots, garlic, leeks) to support liver detoxification pathways
 - ↓ **Table Salt and Processed Foods:** Instead use PRL Pink Salt to obtain electrolyte balancing minerals and many trace minerals
- Include 1 tsp – 1 Tbsp of good quality essential fats/oils** at each meal. Choose cold-pressed, extra virgin olive, coconut, flaxseed and avocado oils, nut butters, and butter made from organic cream. Eliminate any damaged, rancid and hydrogenated oils from the diet such as those found in margarine, fried and take-away food, and many snack foods and baked goods

Recommended Supplementation for John (for approximately 3 months)

BIO EI Multi-trace no Cu: Provides trace minerals and higher doses of essential minerals (B, Se, Mn, Mo, Cr, Zn, Cl and Mg) for balanced mineral levels and natural detoxification processes. Take 1 ml once daily in a little water or juice, after food

BIO EI Chromium: Essential mineral required for blood sugar regulation, protein and cholesterol synthesis. Take 1ml daily in a little water or juice

BIO EI Selenium: Essential mineral for liver function, detoxification of toxic metals, digestive, immune and antioxidant functions in the body. Take 1ml daily in a little water or juice

BIO Elite Ion-Power: Supports proper hydration, sodium and potassium balance, and desirable pH levels. Take 2.5 ml daily added to water bottle and sipped throughout day

BIO Coral Cal-Mag: Supplies easily absorbed and bioavailable Calcium, Magnesium and trace minerals. Helps maintain pH balance. Take 3 capsules daily

BIO Betaine Digest: To support digestion and absorption of minerals. Take 1 capsule with each main meal.

PRL Greens capsules: For phosphorous, calcium and silica and to help balance body pH. Take 2 caps daily with food.

PRL NAC: To replenish glutathione resulting from low sulphur. Helps with toxic metal detoxification. Take 1 capsule twice daily between meals

*All liquids can be taken together in a little juice. We recommend Barkers Blackcurrant (Stevia Sweetened) Cordial to disguise the taste of liquid minerals.

It is recommended to review this protocol with your practitioner monthly to monitor your progress and ensure optimal results. This protocol is intended to build the body up nutritionally in preparation for detoxification of toxic metals.

We strongly recommend you undergo a retest OligoScan in 3 months before commencing a detoxification program under the guidance of a qualified practitioner.

Disclaimer: The information and recommendations provided in this report are not intended to replace the advice of your GP or health care provider. No claims are made in this report to diagnose, treat, cure, or prevent any disease. If symptoms occur, please consult your health care provider to determine the best approach and appropriate modifications that may be required for you.